

Award Number: W81XWH-12-1-0569

TITLE: Functional Analysis of Variants of Unknown Significance in BRCA1 and BRCA2 Using
Complementation of a Synthetic Lethal Interaction with PARP Inhibition

PRINCIPAL INVESTIGATOR: Robert L. Nussbaum, MD

CONTRACTING ORGANIZATION: University of California San Francisco
San Francisco, CA 94103

REPORT DATE: December 2014

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE December 2014		2. REPORT TYPE Final		3. DATES COVERED 30 Sep 2012 - 29 Sep 2014	
4. TITLE AND SUBTITLE Functional Analysis of Variants of Unknown Significance in BRCA1 and BRCA2 Using Complementation of a Synthetic Lethal Interaction with PARP Inhibition				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-12-1-0569	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Robert L. Nussbaum, MD Email: nussbaumr@humgen.ucsf.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of California San Francisco San Francisco, CA 94103				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT We continued to test MCF10A cells containing an inducible shRNA directed against the 3-UTR of BRCA2 for synthetic lethality with PARP inhibitors. Antibody staining with RAD51 and H2AX was used to look for double strand breaks and stalled replication forks and showed increase in RAD51 and H2AX positivity with knock-down of BRCA2 and PARP inhibitors. We also tried very hard to increase transfection efficiency in order to test wild-type and mutant forms of BRCA2 but were not able to transfect the large BRCA2 construct into these cells. We tried lipofection of various types, electroporation, and magnetofection. None of these methods were successful.					
15. SUBJECT TERMS- Variants of Uncertain Significance, BRCA2, Synthetic Lethality, Parp inhibitors					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 9	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code)

TABLE OF CONTENTS

DESCRIPTION	Page
FRONT COVER	1
REPORT DOCUMENTATION PAGE	2
TABLE OF CONTENTS	4
INTRODUCTION	5-6
PROGRESS REPORT BODY	6-8
KEY RESEARCH ACCOMPLISHMENTS	8
REPORTABLE OUTCOMES	8
CONCLUSION	9
REFERENCES	10

INTRODUCTION

An estimated ~22,000 women will be diagnosed with and ~14,000 women will die of **cancer of the ovary** in the United States in 2010¹; approximately 10% of these cases will occur in heterozygotes for a hereditary breast and ovarian cancer syndrome (HBOC)¹ caused by deleterious *BRCA1* or *BRCA2* mutations². Heterozygotes for *BRCA1* or *BRCA2* mutations have a 40% and 11-18% lifetime risk for ovarian cancer respectively, which is far greater than the 1.4% lifetime risk in the general population³⁻⁵. A pathogenic mutation in *BRCA1* or *BRCA2* is an important genetic biomarker for a high ovarian cancer risk in breast cancer patients and their unaffected female relatives; unfortunately 35% and 50% of all DNA variants detected in *BRCA1* and *BRCA2* respectively are variants whose pathogenicity is unknown and are termed “variants of unknown significance” (VUSs)⁶. These missense mutations of unknown significance are located throughout the entire coding sequence of the genes. **Without knowing if patients in whom a VUS is found really have a pathogenic *BRCA1/2* mutation, physicians do not know what to offer these patients and their relatives with regards to risk-reducing surgical procedures or surveillance.**

The functions of the *BRCA1* and *BRCA2* gene products are not completely known but they clearly operate in the homologous recombination (HR) pathway involved in double-strand break repair in DNA^{7,8}. Poly [ADP-ribose] polymerase isoform 1 (Parp1) and its paralog Parp2 also have numerous functions in the cell but, in this context, it is the role of Parp1 in single-strand DNA break (SSB) repair that is probably most important⁹⁻¹³. It is estimated that 10⁴ SSBs normally occur each day in a cell¹⁴. Pharmacological inhibition of Parp1 and Parp2, or siRNA knock-down of Parp1 alone¹⁵, causes a marked increase in SSBs. When a replication fork encounters such SSBs, they become DSBs, which can be detected as foci of phosphorylated γ -H2AX by immunostaining¹⁶. If not repaired, DSBs will cause collapse of the replication fork, leading to apoptosis. The cell normally avoids this catastrophic outcome by repairing the DSBs by homologous recombination, which requires the *BRCA1/2* gene products and can be visualized as foci of RAD51 by immunocytological staining¹⁶. This collaboration between Parp1 and *BRCA1/2* in DNA repair underlies the recently described synthetic lethal interaction (SLI) between them, a synergistic decrease in cell survival in cells with loss of function of either *BRCA1* or *BRCA2* and loss or inhibition of Parp1 activity^{15,17}. SLI has generated intense interest as a new model for chemotherapy and a number of clinical trials of Parp inhibition for breast and ovarian cancers deficient in *BRCA1* or *BRCA2* are underway^{18,19}.

Hypothesis: Missense variants in *BRCA1* and *BRCA2* of unknown pathogenicity identified in cancer patients suspected of having an HBOC syndrome can be reclassified as being either functional (and therefore benign) or non-functional (and therefore pathogenic) based on their ability to complement the SLI caused by inhibiting Parp in cell lines deficient in *BRCA1* or *BRCA2* respectively.

Objective: Develop a robust assay by which individual VUSs can be tested for their effect on *BRCA1/2* function by expressing a *BRCA1* or *BRCA2* gene (cDNA) carrying the VUS into cell lines deficient in *BRCA1* or *BRCA2* respectively, exposing the cells to potent Parp

¹ Abbreviations: HBOC - Hereditary breast and ovarian cancer; *BRCA1/2* – Used when referring to both the *BRCA1* and *BRCA2* genes; VUS – Variant of unknown significance; BIC - Breast Information Core database; SLI -Synthetic lethal interaction; Parp - Poly [ADP-ribose] polymerase; HR – Homologous recombination; SSB – single-strand break in DNA; DSB – Double-strand break in DNA; ORF – Open reading frame; FACS- fluorescence activated cell sorting.

inhibitors, and determining whether the transfected gene is capable of protecting the cell from a SLI, leading to impaired cellular metabolism and loss of viability. **With better information in hand as to the functional significance of VUSs in *BRCA1* and *BRCA2*, patients and their physicians can assess ovarian cancer risk and make much better choices about surveillance and risk-reducing surgery and, ultimately, to more effectively prevent ovarian cancer in these potentially high risk patients.**

BODY OF REPORT


We have accomplished the following Aims of our grant proposal:

SOW Aim 1.

Creation of BRCA2 Knock-Down Cells using shRNA

As described in previous Progress Report, we found two shRNA constructs that in 293T cells were capable of knocking down BRCA2 expression by >70%. We selected one construct, #13 and used a lentivirus vector with a tet-inducible promoter to transfect the MCF10a cell line. Because the lentivirus also expressed an mCherry fluorescent protein, we used flow-sorting to pick the cells with the greatest fluorescence in order to have cell lines with the highest expression.

We used quantitative reverse-transcriptase PCR (qRT-PCR) to measure *BRCA2* RNA with and without doxycycline in the cells infected with shRNA lentivirus as an independent assessment of the effectiveness of knock-down.




Shown here are the cycles for cDNA made from MCF10A cells containing the *BRCA2* shRNA, either without doxycycline or after 24 hours or 72 hours of 100 uM doxycycline. We used *BRCA2*-specific primers and, for reference, primers specific for DNA polymerase alpha. The cDNA was diluted 1/12 and then with two more two-fold dilutions (1/24 and 1/48). The Ct values are very linear, with $r^2 > 0.999$ for all but the *BRCA2* qPCR after 24 hours because of a single point at the 1/24 dilution.

Using DNA polymerase alpha to calculate $\Delta\Delta Ct$ for *BRCA2* shows 24 hours of doxycycline caused a small reduction in *BRCA2* mRNA, but 72 hours caused an increase in $\Delta\Delta Ct$ of ~ 1 , which translates to a 50% drop in mRNA. Recognizing that shRNAs work both by destabilizing mRNA as well as blocking translation, this degree of mRNA reduction is very consistent with a 70% reduction in protein seen by Western blotting.


Synthetic Lethality between Poly (ADP-Ribose) Polymerase (PARP) Inhibition and Knock-Down of *BRCA2*

We had previously established a PARP inhibitor (PARPi) dosage of 10-20 micromolar as being effective in showing synthetic lethality as measured by colony forming assay. We then measured RAD51 and r-H2AX staining of nuclear foci to measure the degree to which doxycycline with or without PARPi increased double-strand breaks. Figure below shows increased intensity of staining when both doxycycline and PARPi are applied to MCF10a cells (Panel C) compared to when either knock-down alone (Panel B) or PARP inhibition alone (Panel A) is used. A summary of all of the experiments are shown in the accompany bar graph in panel D. There is a statistically significant trend of increased RAD51 intensity with both doxycycline and PARPi compared to either alone, although one set of data points, with 50 uM PARPi showed a small drop in the presence of doxycycline compared to without doxycycline.



A summary of staining intensity for RH2AX with or without *BRCA2* knockdown and with or without PARP inhibition is shown below.

RH2AX expression in MCF10a



In the case of RH2AX expression, there is increasing intensity (measured in arbitrary units) when both a knock-down using doxycycline induction of the shRNA and PARP inhibition is present.

Transfection of BRCA2 into cells carrying inducible knock-down of BRCA2

We determined that magnetofection was the least toxic and damaging method of obtaining high frequency transformation of MCF10a cells with, for example, a Green-Fluorescent-Protein expressing vector. **Unfortunately, we could not say the same when we attempted to transfect full-length wildtype BRCA2 expression plasmid into MCF10a cells.** No matter what method of transfection we used, whether it was lipofection, electroporation, or magnetofection, we could not obtain sufficient numbers of transformed cells to be able to use a complementation assay for BRCA constructs carrying missense mutations of unknown functional consequences. We suspect there are two reasons for this: (1) The sheer size of the BRCA2 construct, 16.814 kb, is already very large and difficult to transfect; (2) overexpression of *BRCA2* at certain stages of the cell cycle is very toxic to cells (personal communication from Prof. Alan Ashworth). As a result, we could not transfect many cells and, of those we did, without tight control of expression levels of the protein, we caused unacceptable toxicity in successfully transfected cells.

KEY RESEARCH ACCOMPLISHMENTS

- Development of two cell lines, a highly transformed 293T cell and a less-transformed epithelial cell line, MCF10A, each permanently infected with a lentivirus in which an shRNA against *BRCA2* has been inserted under control of a tetracycline inducible promoter. Treatment of these cells with doxycycline knocks down protein expression of *BRCA2* by >70% by Western blotting and *BRCA2* RNA by ~50% by qPCR.
- Demonstration that treatment of these cells with doxycycline induces cytological signs of synthetic lethality with Parp inhibitor by RAD51 and RH2AX focus formation.

REPORTABLE OUTCOMES

None

CONCLUSION

We have confirmed that there is synthetic lethality for knock-down of BRCA2 and Parp inhibitors in breast epithelial cell line MCF10a, as determined by colony formation, and by RAD51 and RH2AX focus formation.

REFERENCES

1. SEER Incidence and Mortality in Ovarian Cancer. 2008;
<http://seer.cancer.gov/statfacts/html/ovary.html> - incidence-mortality.
2. Risch HA, McLaughlin JR, Cole DE, et al. Prevalence and penetrance of germline *BRCA1* and *BRCA2* mutations in a population series of 649 women with ovarian cancer. *American journal of human genetics*. Mar 2001;68(3):700-710.
3. Chen S, Parmigiani G. Meta-analysis of *BRCA1* and *BRCA2* penetrance. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. Apr 10 2007;25(11):1329-1333.
4. Chen S, Iversen ES, Friebel T, et al. Characterization of *BRCA1* and *BRCA2* mutations in a large United States sample. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. Feb 20 2006;24(6):863-871.
5. Antoniou A, Pharoah PD, Narod S, et al. Average risks of breast and ovarian cancer associated with *BRCA1* or *BRCA2* mutations detected in case Series unselected for family history: a combined analysis of 22 studies. *American journal of human genetics*. May 2003;72(5):1117-1130.
6. Carvalho MA, Marsillac SM, Karchin R, et al. Determination of cancer risk associated with germ line *BRCA1* missense variants by functional analysis. *Cancer Res*. Feb 15 2007;67(4):1494-1501.
7. Evers B, Helleday T, Jonkers J. Targeting homologous recombination repair defects in cancer. *Trends Pharmacol Sci*. Aug 2010;31(8):372-380.
8. Gudmundsdottir K, Ashworth A. The roles of *BRCA1* and *BRCA2* and associated proteins in the maintenance of genomic stability. *Oncogene*. Sep 25 2006;25(43):5864-5874.
9. Megnin-Chanet F, Bollet MA, Hall J. Targeting poly(ADP-ribose) polymerase activity for cancer therapy. *Cell Mol Life Sci*. Nov 2010;67(21):3649-3662.
10. Carden CP, Yap TA, Kaye SB. PARP inhibition: targeting the Achilles' heel of DNA repair to treat germline and sporadic ovarian cancers. *Curr Opin Oncol*. Sep 2010;22(5):473-480.
11. Helleday T, Bryant HE, Schultz N. Poly(ADP-ribose) polymerase (PARP-1) in homologous recombination and as a target for cancer therapy. *Cell Cycle*. Sep 2005;4(9):1176-1178.
12. Dantzer F, Schreiber V, Niedergang C, et al. Involvement of poly(ADP-ribose) polymerase in base excision repair. *Biochimie*. Jan-Feb 1999;81(1-2):69-75.
13. Dantzer F, de La Rubia G, Menissier-De Murcia J, Hostomsky Z, de Murcia G, Schreiber V. Base excision repair is impaired in mammalian cells lacking Poly(ADP-ribose) polymerase-1. *Biochemistry*. Jun 27 2000;39(25):7559-7569.
14. Lindahl T. Instability and decay of the primary structure of DNA. *Nature*. Apr 22 1993;362(6422):709-715.
15. Bryant HE, Schultz N, Thomas HD, et al. Specific killing of *BRCA2*-deficient tumours with inhibitors of poly(ADP-ribose) polymerase. *Nature*. Apr 14 2005;434(7035):913-917.
16. Saleh-Gohari N, Bryant HE, Schultz N, Parker KM, Cassel TN, Helleday T. Spontaneous homologous recombination is induced by collapsed replication forks that are caused by endogenous DNA single-strand breaks. *Molecular and cellular biology*. Aug 2005;25(16):7158-7169.
17. Farmer H, McCabe N, Lord CJ, et al. Targeting the DNA repair defect in BRCA mutant cells as a therapeutic strategy. *Nature*. Apr 14 2005;434(7035):917-921.
18. Boss DS, Beijnen JH, Schellens JH. Inducing synthetic lethality using PARP inhibitors. *Curr Clin Pharmacol*. Aug 2010;5(3):192-195.
19. Iglehart JD, Silver DP. Synthetic lethality--a new direction in cancer-drug development. *N Engl J Med*. Jul 9 2009;361(2):189-191.